

Editorial comment 179

A rare 19th century French obstetrical atlas 180

HISTORICAL REVIEW

Vaginal birth after classical Caesarean section 182
de Costa C

This review examines the lack of evidence to support the current belief that the scar of a classical Caesarean section is more prone to rupture than that of a lower segment Caesarean section.

ORIGINAL ARTICLES

Paravaginal defects: a comparison of clinical examination and 2D/3D ultrasound imaging 187
Dietz HP, Pang S, Korda A, Benness BC

This prospective clinical and ultrasound study of 59 women without previous prolapse or incontinence surgery shows that pelvic floor ultrasound does not correlate well with clinical assessment for paravaginal defects.

Transperineal versus transvaginal ultrasonographic evaluation of the cervix at each trimester in normal pregnant women 191
Ozdemir I, Demirci F, Yucel O

This study of normal pregnant women suggests that transperineal ultrasonography is a satisfactory alternative to a transvaginal evaluation of the cervix throughout pregnancy.

Bowel, bladder and sexual function in women undergoing laparoscopic posterior compartment repair in the presence of apical or anterior compartment dysfunction 195

Thornton MJ, Lam A, King DW

This prospective study of the functional outcome of laparoscopic posterior compartment repair in the presence of anterior compartment dysfunction shows that anatomical improvement may not be followed by long-term functional improvement.

Management of preterm prelabour rupture of membranes: an audit. How do the results compare with clinical practice guidelines? 201

Giles M, Garland S, Oats JJN

This retrospective audit of the introduction of a new clinical practice guideline shows overall good compliance, although antibiotic prescribing and management of Group B streptococcus colonization could be improved, and an element of the guideline was seen to not be useful.

Antenatal antecedents of moderate or severe neonatal encephalopathy in term infants – a regional review 207

West CR, Curr L, Battin MR, Harding JE, McCowan LM, Belgrave S, Knight DB, Westgate JA

This audit of term infants admitted with moderate-severe neonatal encephalopathy shows that peripartum events were the major contributors to neurological damage in these infants.

Presenting symptoms of epithelial ovarian cancer 211

Lataifeh I, Marsden DE, Robertson G, Gebiski V, Hacker NF

This retrospective cohort study comparing women with stage 1 and stage 3 epithelial ovarian cancer shows that women presenting with early disease appear to be younger and have biologically different disease when compared to those with advanced disease.

Comparison of a self-administered tampon ThinPrep test with conventional pap smears for cervical cytology 215

Budge M, Halford J, Haran M, Mein J, Wright G

This is a comparative observational study of self-administered tampon specimen as an alternative method of detecting cytological abnormalities to the conventional Papanicolaou smear.

Substance use during pregnancy: risk factors and obstetric and perinatal outcomes in South Australia 220

Kennare R, Heard A, Chan A

Substance use by 707 women, of 89,080 confinements, was associated with increased risks for antepartum haemorrhage and poor perinatal outcomes; however, only a small amount of the variance in outcomes can be explained by the substance use alone.

Physiotherapy for female stress urinary incontinence: a multi-centre observational study 226

Neumann PB, Grimmer KA, Grant RE, Gill VA

This prospective multicentre clinical study of physiotherapy management of female stress urinary incontinence in 208 women shows it to be an effective treatment option.

SHORT COMMUNICATIONS

Primary post-partum haemorrhage in an Australian tertiary hospital: a case-control study 233

Henry A, Birch M-R, Sullivan EA, Katz S, Wang YA

This case-control study of 250 women delivering vaginally at a tertiary hospital examines the 1 in 8 women having a postpartum haemorrhage, and confirms risk factors to be a past history of postpartum haemorrhage, a long second stage of labour, assisted vaginal birth, and incomplete placenta/ragged membranes.

Use of urethral plugs for urinary incontinence following fistula repair 237

Goh JTW, Browning A

This manuscript describes a two-case pilot study of the use of urethral plugs as an effective short-term management of women with post-fistula incontinence.

CASE SERIES

Outcomes of pregnancy complicated by thyroid disease 239

Harborne LR, Alexander CE, Thomson AJ, O'Reilly DS, Greer IA

This retrospective review of 81 ongoing pregnancies in women with a history of thyroid dysfunction shows that pregnancy outcome was good in the majority of cases given appropriate replacement therapy for stated reference values.

Contents 2

Pregnancy and the Rastelli operation 243
Radford DJ, Stafford G
This case series of six women who were survivors of the Rastelli repair operation for complex congenital heart disease and subsequently became pregnant shows that successful pregnancy is possible, but careful cardiac evaluation before, during and after pregnancy is essential.

LETTERS TO THE EDITOR

CLINICAL-SCIENTIFIC NOTES

Unexpected complication of uterine myoma morcellation 248
Rekha W, Amita M, Sudeep G, Hemant T
This case report highlights the risk of intra-peritoneal dissemination of an unsuspected leiomyosarcoma in the event of a morcellator being used for retrieval of the specimen at laparoscopic surgery.

Sheehan's syndrome presenting post-partum with diabetes insipidus 249
Weston G, Chaves N, Bowditch J
Though Sheehan's syndrome usually affects only the anterior pituitary, this unusual case reports infarction of the posterior pituitary, manifesting as diabetes insipidus.

A new surgical technique for dealing with uterine inversion 250
Robson S, Adair S, Bland P
The authors describe a new surgical technique for abdominal correction of uterine inversion.

Pityriasis Rosea in pregnancy – specific diagnostic implications and management considerations 252
Chuh AAT, Lee A, Chan PKS
The authors describe two women with pityriasis rosea complicating pregnancy, and discuss its important differential diagnosis.

Isolated vulval varicosity in the non-pregnant state: a case report with review of the treatment options 254
Leung SW, Leung PL, Yuen PM, Rogers, MS
This case report describes the use of laparoscopic ovarian vein ligation for an isolated vulval varicosity in a non-pregnant lady.

Life-threatening uterine haemorrhage six weeks after Caesarean section due to uterine scar endometriosis: Case report and review of literature 256
Sholapurkar SL, Sharp NC, Hirschowitz L
This unusual case report describes severe haemorrhage from uterine wall caesarean section scar endometriosis only 6 weeks after the initial operation.

CORRESPONDENCE 259