



C-Gyn 3

Intrauterine contraceptive devices and infection

The Royal Australian and New Zealand College of Obstetricians and Gynaecologists (RANZCOG) endorses the views of the International Planned Parenthood Federation (IPPF) that the intrauterine contraceptive device (IUCD) is an effective and safe method of contraception for properly screened women. The IPPF has published a detailed statement about the use of IUCDs, which is a useful resource.

Currently available devices include the copper containing Cu T380 and Multiload Cu 375, as well as the levonorgestrel containing Mirena, which may be used to treat menorrhagia as well as offering contraception.

A careful history and examination is essential to identify any relative or absolute contraindications to the use of an Intrauterine Contraceptive Device (IUCD). Sexually transmitted infection (STI) risk should be assessed and where appropriate testing for chlamydia trachomatis, neisseria gonorrhoea and other STI's as requested by the woman be considered prior to insertion or change of an IUCD. Ideally these results should be available prior to IUCD insertion.

All patients should be counselled about the effectiveness and failure rates of IUCDs and their possible short and long-term complications, including menstrual changes and pelvic infection.

Both oral and written information should be provided. At the time of insertion of an IUCD any possibility of pregnancy should be excluded.

World Health Organization (WHO) studies have demonstrated a small risk of pelvic infection in the first 20 days after insertion, often relating to asymptomatic and unrecognised STIs. Following this period of increased risk the rate of Pelvic Inflammatory Disease (PID) in IUCD users is approximately the same as would be expected in the general population not using IUCDs.

The balance of evidence suggests that the use of an IUCD does not affect return to fertility in nulliparous or multiparous women.

Women at increased risk of STI acquisition should be counselled about other methods of contraception - WHO recommends that the risks generally outweigh the benefits in women at increased risk of STI's. This risk clearly relates to the sexual relationship/s of the woman and her partner/s but is greater on a population basis for women younger than 25 years of age. FFPRHC (Faculty of Family Planning and Reproductive Health Care RCOG) states that women at increased risk of STI's after counseling may still choose an IUCD and safe sex and condom use should be promoted.

Patients should be advised to attend for review 4 to 6 weeks after insertion of the IUCD and thereafter at least every 1-2 years. They should also be advised to present if abnormal bleeding, or symptoms suggestive of infection or pregnancy occur, or if they are unable to locate the string of the device. Information should be provided advising when renewal of the device is required.

Actinomyces

Pap smears in asymptomatic women are performed every two years in Australia and every three years in New Zealand. Cytological examination may reveal Actinomyces-like organisms (ALOs). It is important to be aware that some ALOs reported in Pap smears are not in fact Actinomyces or are Actinomyces species which do not commonly cause sepsis. Furthermore the presence of ALOs in a Pap smear does not correlate well with culture for Actinomyces nor with the risk of subsequent PID. Certainly the presence of ALOs in the Pap smear of a woman who is asymptomatic should not alone be considered an indication for removal of the IUCD.

Recommendations for management vary. A suggested course of action would be to take a cervical swab for specific formal culture of Actinomyces, which may require a specialized laboratory, special media and/or prior notification of the laboratory. IUCD use may continue provided the culture is negative and the woman asymptomatic. Identification of Actinomyces israelii on culture, or direct immuno-fluorescence if available, should always indicate removal of an IUCD; if the patient is symptomatic prolonged anti-microbial treatment should be used in consultation with a clinical microbiologist or infectious diseases physician; surgery may need to be considered to drain any associated collections.

Pregnancy with an IUCD in situ

If pregnancy is diagnosed with an IUCD in situ, the device should be removed if possible because of the risk of serious complications. These complications include a 50% incidence of spontaneous miscarriage and an increased incidence of placental problems such as antepartum haemorrhage, threatened premature labour and adherent placenta.

References

1. Arend S, Oosterhoof H, van Dissel JT. Actinomyces and the intrauterine device. Archives of Internal Medicine 1998;138:1270.
2. Farley TMM, Rowe PJ, Rosenberg MJ, Chen J-H, O M. Intrauterine devices and pelvic inflammatory disease: an international perspective. Lancet 1992;339:785-788.
3. IPPF. International Medical Advisory Panel statement on intrauterine devices. IPPF Medical Bulletin 1995;29(6).
<http://www.ippf.org/en/Resources/Medical/IMAP+statements.htm>
4. ACOG Practice Bulletin No. 59. Intrauterine Device. Obstetrics and Gynaecology 2005; 105 (1): 223-32.
5. RCOG FFPRHC Clinical guidance Intrauterine Contraception CEU Nov 2007 ISSN1755-103X
<http://www.ffprhc.org.uk/default.asp?Section=GoodMedicalSubsection=ClinicalGuidance>
6. World Health Organisation (WHO) Medical Eligibility Criteria for contraceptive use. Geneva Switzerland: WHO, 3rd edn 2004.
7. National Institute for Health and Clinical Guidance(NICE) long acting reversible contraception: the effective and appropriate use of long acting reversible contraception 2005 <http://www.nice.org.uk/pdf/CG030fullguidance.pdf>

Disclaimer

This College Statement is intended to provide general advice to Practitioners. The statement should never be relied on as a substitute for proper assessment with respect to the particular circumstances of each case and the needs of each patient.

The statement has been prepared having regard to general circumstances. It is the responsibility of each Practitioner to have regard to the particular circumstances of each case, and the application of this statement in each case. In particular, clinical management must always be responsive to the needs of the individual patient and the particular circumstances of each case.

This College statement has been prepared having regard to the information available at the time of its preparation, and each Practitioner must have regard to relevant information, research or material which may have been published or become available subsequently.

Whilst the College endeavours to ensure that College statements are accurate and current at the time of their preparation, it takes no responsibility for matters arising from changed circumstances or information or material that may have become available after the date of the statements.