



## **College Statement**

---

Title	<b>Umbilical Cord Blood Banking</b>
Statement No.	<b>C-Obs 18</b>
Date of this document	<b>July 2007</b>
First endorsed by Council	<b>February 2003</b>
Next review due:	<b>November 2009</b>

---

### **Introduction**

Umbilical Cord blood represents a rich source of Haemopoetic Stem Cells (HSC). The major clinical use of cord blood stem cells has been in the treatment of haematological malignancy in children. Currently about 20% of stem cell transplants for young patients are cord blood transplants. Cord blood as an alternative to bone marrow as a source of HSC for allogeneic transplantation has a number of potential advantages for both adults and children including availability, extension of the donor pool, lower incidence of graft versus host disease and lower incidence of viral transmission.

### **Current uses of cord blood include**

1. *Cord blood transplants (CBT) from related donors*  
Cord blood transplantation is particularly useful for treating children with haemoglobinopathies
2. *Cord blood transplants in children from unrelated donors*  
CBT is a good option for children with haematological malignancy who lack a related donor.
3. *Cord blood transplants in adults from unrelated donors*  
Results of HSC transplants in adults with haematological malignancy are encouraging

There is substantial speculation about the use of cord blood non-HSC in treatment of a variety of acute and chronic conditions and there is increasing interest in the use of fetal-derived stem cells in the treatment of neurological and other disorders.

In 2000, the Commonwealth Government assisted in the establishment of an Australian National Cord Blood Collection Network (NCBCN). The Australian Bone Marrow Donor Registry (ABMDR) was contracted to manage the (four year) "Development Phase" of the National Cord Blood Collection Network.

The network was officially launched under the name “AusCord” in September 2002.

The main objectives of the network is to collect, process and store 22,000 searchable, TGA compliant Cord Blood Units, (including 2,000 indigenous units) for the purpose of transplantation and to establish a computerised national registry of Cord Blood Units for the purpose of searching, matching and distributing Cord Blood Units.

## **Types of Cord Blood Donation**

### *4. Altruistic (non directed) donations*

Thirty percent (30 percent) of Australian patients in need of an allogeneic bone marrow transplant have a suitable family donor, that is, 5 or 6 out of 6 Human Leucocyte Antigen (HLA) matched close relative. Of the remaining 70 percent of patients, only 20 – 25 percent are able to find an unrelated bone marrow donor on the existing registries. Therefore, over 50 percent of patients in need of an allogeneic transplant do not have a histocompatible related or unrelated donor. This same group may be eligible for a cord blood transplant as it is known that the degree of compatibility between the patient and the cord unit does not need to be as stringent as in bone marrow transplantation, with matching acceptable to a level of 4 out of 6 match. This is because graft-versus-host disease is lower in cord blood transplants at a given level of match, thereby increasing an individual's chances of finding a suitable cord blood donation at an acceptable level of match. The main indication for cord blood transplantation is relapsed acute lymphoblastic leukaemia in children.

Following the experience at the New York Cord Blood Bank and the National Marrow Donor Program in the United States in identifying suitably matched donors (5 out of 6 and 6 out of 6 HLA matches) for patients, the NCBCN expect that a collection of 20,000 cord blood units will enable 80 – 90% of Australian requests to be met.

### *5. Directed donations in at risk families*

The collection of cord blood units for use by siblings born into a family where there is a known genetic disease amenable to HSC transplant remains a recommendation. If the cells are HLA-compatible they may be used for an affected child. If not, they may be useable for a future HLA-compatible sibling.

### *6. Directed donations in low-risk families*

It is estimated that of approximately 200,000 placental blood units stored for the exclusive use of the baby from whose birth the placental blood was obtained, at most, 74 units (and this is very likely a gross overestimate) might “possibly” be used and 199,926 would be taken out of circulation for use by patients who require an allogeneic transplant <sup>1</sup>.

This is a rapidly progressive area of clinical medicine. It is possible that future advances may attach considerable value to these cells, in the event of childhood or adult-onset disease in the individual from whom the stem cells were obtained.

On the other hand, it is also possible that parallel advances in genetic manipulation of donor stem cells may render the expense of a directed donation unnecessary.

If a donor developed a haematological malignancy and required a transplant in the future their own stored cord blood is generally not recommended to treat their subsequent malignancy. The use of that cord blood for an HLA-compatible relative is theoretically acceptable if permitted in the contract with the private facility storing the stem cells.

At this stage, each individual should make a decision, weighing the considerable cost of directed donation against an uncertain probability of future benefit.

## **Practical implications of cord blood collection**

A limited number of hospitals are funded to arrange altruistic cord blood collections for AusCord.

Cord blood collection for private storage does impose some burden on obstetricians and midwives, including:

- Completion of associated paperwork.
- Interference in the management of the third stage to allow cord blood collection at a time when the mother is at risk of postpartum haemorrhage.
- Pressure to collect sufficiently large volume of blood as there is a minimum requirement in the volume of blood that can be stored.
- The requirement for adequate aseptic technique.
- The necessity for staff in public hospitals to be trained to perform cord blood collection.

Hospitals need to develop their own policy on how to respond to prenatal request for cord blood storage through commercial providers.

## **Recommendations**

The Royal Australian and New Zealand College of Obstetricians and Gynaecologists (RANZCOG) supports the collection of altruistic and directed cord blood donations in at risk families.

The routine collection of directed donations in low-risk families must consider both the cost and an uncertain probability of future benefit.

## **References**

1. Placental Blood Transplant and Autologous Banking – Caveat Emptor Johnson, F.L. *Journal of Paediatric Haematology / Oncology*, Volume 19 (3); 183 – 186.

## **Disclaimer**

This College Statement is intended to provide general advice to Practitioners. The statement should never be relied on as a substitute for proper assessment with respect to the particular circumstances of each case and the needs of each patient.

The statement has been prepared having regard to general circumstances. It is the responsibility of each Practitioner to have regard to the particular circumstances of each case, and the application of this statement in each case. In particular, clinical management must always be responsive to the needs of the individual patient and the particular circumstances of each case.

This College statement has been prepared having regard to the information available at the time of its preparation, and each Practitioner must have regard to relevant information, research or material which may have been published or become available subsequently.

Whilst the College endeavours to ensure that College statements are accurate and current at the time of their preparation, it takes no responsibility for matters arising from changed circumstances or information or material that may have become available after the date of the statements.