



## C-Obs 7

# Diagnosis of Gestational Diabetes Mellitus

Current evidence suggests that there is a benefit in reduced perinatal morbidity in screening for and treating gestational diabetes mellitus. When screening for gestational diabetes, there should be uniformity in the testing used and the subsequent follow up. The following steps are recommended:

Biochemical screening for gestational diabetes should be performed between 26 and 28 weeks of gestation. Earlier testing may be performed in women at particularly high risk of gestational diabetes (e.g. gestational diabetes in a previous pregnancy).

Most clinicians will choose to recommend a Glucose Challenge Test (GCT) and then request a Glucose Tolerance Test (GTT) if the GCT is abnormal. In a GCT, plasma glucose should be measured 1 hour after either a 50g or 75g oral glucose load given in the non-fasting state. Women with a plasma glucose value at 1 hour after a 50g load of  $\geq 7.8$  mmol/l or after a 75g load of  $\geq 8.0$  mmol/l are regarded as positive and should undergo a full glucose tolerance test, (fasting, 75g glucose, 0, and 2 hr glucose estimations). In addition to the above, women at particularly high risk of gestational diabetes, such as women who had gestational diabetes in a previous pregnancy, may be tested earlier in pregnancy.

Some clinicians may choose to omit the GCT and recommend a full GTT. They may so recommend for all pregnant women or only those with a high likelihood of recall after a GCT. These clinicians may feel that a dual testing regimen (GCT then GTT if GCT abnormal) is inconvenient for the women involved, increases the administrative workload (recall processes) and inevitably delays the commencement of therapy.

The diagnosis of gestational diabetes should be made on a GTT following a fasting, 75g 2hr glucose tolerance test. Fasting glucose level of  $\geq 5.5$ mmol/l is diagnostic of gestational diabetes, as is a 2 hr level  $\geq 8.0$  mmol/l by Australian criteria, or  $\geq 9.0$  mmol/l by NZ criteria.

These recommendations must not preclude confirmation of diabetes at any stage of pregnancy if there are clinical features to suggest the diagnosis.

Women who had gestational diabetes should have a repeat 75g oral glucose tolerance test performed 6 to 8 weeks after delivery. This should be evaluated according to standard WHO criteria for the non-pregnant state. Women who do not have diabetes mellitus at this time should still be regarded as at risk of developing diabetes mellitus later in life and should be screened every two to three years.

## References

1. Revision of guidelines for the management of gestational diabetes mellitus, Letter to the Editor, J J N Oats and H D McIntyre, MJA September 2004;181(6): 342  
[http://www.mja.com.au/public/issues/181\\_06\\_200904/letters\\_200904\\_fm-2.html](http://www.mja.com.au/public/issues/181_06_200904/letters_200904_fm-2.html)

2. [Gestational diabetes mellitus – management guidelines](#), updated by ADIPS December 2002, (amendment published as a letter to the Editor of the MJA).
3. ADIPS Gestational diabetes mellitus – management guidelines, L Hoffman, C Nolan, J D Wilson, J J N Oats and D Simmons, MJA 1998; 169: 93-97  
<http://www.mja.com.au/public/issues/jul20/hoffman/hoffman.html>
4. Crowther CA, Hiller JE, Moss, JR, McPhee AJ, Jeffries WS and Robinson JS: Australian Carbohydrate Intolerance Study in Pregnant Women (ACHOIS) Trial Group. Effect of Treatment of Gestational Diabetes Mellitus on Pregnancy Outcomes. The New England Journal of Medicine. 2005; 352(24):2477-2486. <http://content.nejm.org/>

## Links to other related College Statements

### [C-Obs 3 Antenatal screening tests](#)

#### **Disclaimer**

This College Statement is intended to provide general advice to Practitioners. The statement should never be relied on as a substitute for proper assessment with respect to the particular circumstances of each case and the needs of each patient.

The statement has been prepared having regard to general circumstances. It is the responsibility of each Practitioner to have regard to the particular circumstances of each case, and the application of this statement in each case. In particular, clinical management must always be responsive to the needs of the individual patient and the particular circumstances of each case.

This College statement has been prepared having regard to the information available at the time of its preparation, and each Practitioner must have regard to relevant information, research or material which may have been published or become available subsequently.

Whilst the College endeavours to ensure that College statements are accurate and current at the time of their preparation, it takes no responsibility for matters arising from changed circumstances or information or material that may have become available after the date of the statements.